



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

BIBLIOGRAPHIC RECORD TARGET

Graduate Library
University of Michigan

Preservation Office

Storage Number: _____

ACE2795

UL FMT B RT a BL m T/C DT 07/18/88 R/DT 07/18/88 CC STAT mm E/L 1

010: : |a 15005645

035/1: : |a (RLIN)MIUG84-B10868

035/2: : |a (CaOTULAS)160253297

040: : |c MiU |d MiU

050/1:0: : |a QA484 |b .B65

100:1 : |a Brennan, Michael H.

245:10: |a Reciprocal arcs, |c by Michael H. Brennan.

260: : |a Lancaster, Pa., |b Press of the New era printing company, |c 1915.

300/1: : |a 7 p. |b diagrs. |c 26 cm.

650/1: 0: |a Circle

998: : |c EM |s 9124

Scanned by Imagenes Digitales
Nogales, AZ


On behalf of
Preservation Division
The University of Michigan Libraries

Date work Began: _____
Camera Operator: _____

RECIPROCAL ARCS

BY

MICHAEL H. BRENNAN



PRESS OF
 THE NEW ERA PRINTING COMPANY
 LANCASTER, PA.

1915

Copyright, 1915
By MICHAEL H. BRENNAN

DEVILS LAKE, N. D., February 9, 1915.

I propose to demonstrate the following proposition:

In a circle whose radius is 1, the distance from the center to the middle point of the chord of any arc, is the square of the chord of one-third of another arc in the circle; and the distance from the center to the middle point of the chord of that other arc, is the square of the chord of one-third of the former arc.

For convenience of construction and demonstration, the chords of the respective arcs are taken perpendicular to the horizontal diameter, so that the cosine of half the arc on one side of the center will be shown to be the square of the chord of one-third of the whole arc on the other side, and conversely.



3

Thus, in Fig. 1, let $HAECK$ be the circle, MN its horizontal diameter produced, AC the chord of the given arc, AE its half, OB the cosine of the arc AE . On DB as a diameter describe a circle cutting the perpendicular diameter at F . Let DO , the radius, equal 1. Then FO is a mean proportional between DO and OB . Hence $OB = FO^2$. Call FO , y , and OB , y^2 . Then $BE = 1 - y^2$.

A circle with F as a center and $BE = 1 - y^2$ as a radius will cut the main circle at H . From H drop HK perpendicular to DE , cutting it at I . On IE as a diameter describe a circle cutting the perpendicular diameter at G . GO will be a mean proportional between IO and OE , and $IO = GO^2$. Call GO , x , and IO , x^2 , then is OB the square of the chord of one-third of the arc HDK , and IO is the square of the chord of one-third of the arc AEC .

Proof: Draw Ft perpendicular to FO and HI , and draw AT perpendicular to FO and AB . As $IO = GO^2 = x^2$, $DI = 1 - x^2$. Through A and G' draw a line cutting the circle at P and the diameter DE produced, at M . Produce HF cutting the circle at Q and the diameter DE produced, at N .

$$AB = \frac{1}{2}AC = \sqrt{1 - OB^2} = \sqrt{1 - y^4}.$$

$GA^2 = TA^2 + TG^2$; but $TA = OB = y^2$, and $TG = AB - GO$. Let $AB = c$. Then $GA = \sqrt{y^4 + (c - x)^2}$, and as $c^2 = 1 - y^4$,




FIG. 2.

$$GA = \sqrt{y^4 + 1 - y^4 + x^2 - 2cx} = \sqrt{x^2 - 2cx + 1}.$$

(See Figure 2.)

For convenience, let $HI = k$, $FN = \alpha$ and $GM = a$. From similar triangles, FNO and HFT , we have $FN : FO :: 1 - y^2 : HI - FO = Ht$, or,

$$\alpha : y :: 1 - y^2 : k - y; \quad y - y^3 = \alpha(k - y); \quad \alpha = (y - y^3)/(k - y) \quad (1)$$

$$(1 - y^2)^2 = \overline{Ft}^2 + \overline{Ht}^2 = (HI - tI)^2 + t\overline{F}^2 = (k - y)^2 + x^4.$$

As $tF = IO = x^2$, $x^4 = 1 - k^2$. Substituting in $(1 - y^2)^2 = (k - y)^2 + x^4$, we have

$$1 - 2y^2 + y^4 = y^2 - 2ky + 1; \quad y^4 - 3y^2 + 2ky = 0; \quad y^3 - 3y + 2k = 0,$$

which may be written

$$y - y^3 = 2k - 2y; \quad (y - y^3) \div (k - y) = 2.$$

We have shown that $(y - y^3) \div (k - y) = \alpha$, whence $\alpha = 2 = FN = DE$, the diameter of the circle. The circumscribing circle of the rightangled triangle FNO will have for a radius one-half of α or of 2, and a radius from the middle point of α to the vertex at O will = 1; but a line drawn to Q where α cuts the circle is equal to 1; hence FN is bisected by the circle at the point Q . Hence angle $QON = QNO$, and $FQO = 2QNO$. $HOD = OHQ + QNO = HQO + QNO$. But $QNO = \frac{1}{2}OQH$; whence $QNO =$ one-third of HOD and arc $QE =$ one-third of arc HD .

In the foregoing the arc HD is found as a result of the construction, but its trisection does not give any measure of the one-third of AE . If a should be found equal to α or if GA is equal to $1 - x^2$, the x equation would have the same form as the y equation, and without further reasoning the mind would conclude that the arc PD is equal to one-third of the arc AE .

Having found that $GA = \sqrt{x^2 - 2cx + 1}$ we have from similar triangles, GMO and GAT , the proportion:

$$GA = \sqrt{x^2 - 2cx + 1} : a :: c - x : x; \quad x\sqrt{x^2 - 2cx + 1} = a(c - x).$$

Squaring, we have:

$$x^4 - 2cx^3 + x^2 = a^2c^2 - 2a^2cx + a^2x^2 = 0,$$

$$x^4 - 2cx^3 + (1 - a^2)x^2 + 2a^2cx - a^2c^2 = 0.$$

Note.—That the only given value we have at this stage in the equation is c , and attention is called to the fact that we are concerned with the value of a only and not of x .

The different operations by which the value of a is arrived at will now be given.

In Fig. 1, LR and PA being chords intersecting at G , the product of the segments of one equals the product of the segments of the other. Let $PG = s$. We have $LG \cdot GR = AG \cdot PG$; $GO = x$, $LG = 1 - x$ and $GR = 1 + x$; AG has been shown to be $\sqrt{x^2 - 2cx + 1}$; whence $1 - x^2 = s\sqrt{x^2 - 2cx + 1}$; $\sqrt{x^2 - 2cx + 1} = (1 - x^2) \div s$, and $s = (1 - x^2) \div \sqrt{x^2 - 2cx + 1}$.

$MG = a$; $GO = x$; $a + \sqrt{x^2 - 2cx + 1} = a + (1 - x^2) \div s$ and $AB = c$. Whence, from similar triangles:

